

Advil PC

Advil® Concentrated Drops

Drug Facts

Active ingredient
(in each 1.25 mL)

Purposes

Ibuprofen 50 mg

Fever reducer/Pain reliever

Uses

temporarily:

reduces fever

relieves minor aches and pains due to the common cold, flu, headaches and toothaches

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction which may include:

hives facial swelling asthma (wheezing) shock

Stomach bleeding warning: Taking more than recommended may cause stomach bleeding.

Do not use if the child has ever had an allergic reaction to any other fever reducer/pain reliever

Ask a doctor before use if the child has

high blood pressure, heart or kidney disease or is taking a diuretic

not been drinking fluids

ulcers

bleeding problems

lost a lot of fluid due to continued vomiting or diarrhea

stomach problems that last or come back, such as heartburn, upset stomach or pain

problems or serious side effects from taking fever reducers or pain relievers

Ask a doctor or pharmacist before use if the child is

under a doctor's care for any serious condition taking any other product that contains

ibuprofen, or any other pain reliever/fever reducer taking a prescription product for

anticoagulation (blood thinning) taking any other drug

When using this product give with food or milk if stomach upset occurs

Stop use and ask a doctor if

an allergic reaction occurs. Seek medical help right away. fever or pain gets worse or lasts

more than 3 days the child does not get any relief within first day (24 hours) of treatment

stomach pain or upset gets worse or lasts redness or swelling is present in the painful area

any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison

Control Center right away.

Directions

do not give more than directed

shake well before using

find right dose on chart below. If possible, use weight to dose; otherwise use age.

repeat dose every 6-8 hours, if needed

do not use more than 4 times a day

measure with the dosing device provided. Do not use with any other device.

Dosing Chart

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lb	6-11 mos	1.25 mL
18-23 lb	12-23 mos	1.875 mL

Other information

one dose lasts 6-8 hours

store at 20-25°C (68-77°F)

see bottom of box for lot number and expiration date

Inactive ingredients

GRAPE FLAVOR

artificial flavor, carboxymethylcellulose sodium, citric acid, edetate disodium, FD&C blue no.1,

FD&C red no. 40, glycerin, microcrystalline cellulose, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

WHITE GRAPE FLAVOR

artificial flavor, carboxymethylcellulose sodium, citric acid, edetate disodium, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

Children's Advil® Suspension

Drug Facts

Active ingredient (in each 5mL)

Purposes

Ibuprofen 100 mg

Fever reducer/Pain reliever

Uses

temporarily:

reduces fever

relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction which may include:

hives facial swelling asthma (wheezing) shock

Stomach bleeding warning: Taking more than recommended may cause stomach bleeding.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use if the child has ever had an allergic reaction to any other fever reducer/pain reliever

Ask a doctor before use if the child has

not been drinking fluids ulcers bleeding problems

lost a lot of fluid due to continued vomiting or diarrhea

problems or serious side effects from taking fever reducers or pain relievers

stomach problems that last or come back, such as heartburn, upset stomach or pain

high blood pressure, heart or kidney disease or is taking a diuretic

Ask a doctor or pharmacist before use if the child is

under a doctor's care for any serious condition taking any other product that contains

ibuprofen, or any other pain reliever/fever reducer taking a prescription product for

anticoagulation (blood thinning) taking any other drug

When using this product give with food or milk if stomach upset occurs

Stop use and ask a doctor if

an allergic reaction occurs. Seek medical help right away. fever or pain gets worse or lasts

more than 3 days the child does not get any relief within first day (24 hours) of treatment

stomach pain or upset gets worse or lasts redness or swelling is present in the painful area

any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison

Control Center right away.

Directions

do not give more than directed

shake well before using

find right dose on chart below. If possible, use weight to dose; otherwise use age.

repeat dose every 6-8 hours, if needed

do not use more than 4 times a day

measure only with the blue dosing cup provided. Blue dosing cup to be used with Children's

Advil Suspension only. Do not use with other products. Dose lines account for product

remaining in cup due to thickness of suspension.

Dosing Chart

Weight (lb)	Age (yr)	Dose (tsp)
under 24 lb	under 2 yr	ask a doctor
24-35 lb	2-3 yr	1 tsp

36-47 lb	4-5 yr	1 1/2 tsp
48-59 lb	6-8 yr	2 tsp
60-71 lb	9-10 yr	2 1/2 tsp
72-95 lb	11 yr	3 tsp

Other information

one dose lasts 6-8 hours

store at 20-25°C (68-77°F)

see side of box for lot number and expiration date

Inactive ingredients

FRUIT FLAVOR

artificial flavors, carboxymethylcellulose sodium, citric acid, edetate disodium, FD&C red no. 40, glycerin, microcrystalline cellulose, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

GRAPE FLAVOR

artificial flavor, carboxymethylcellulose sodium, citric acid, edetate disodium, FD&C blue no.1, FD&C red no. 40, glycerin, microcrystalline cellulose, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

BLUE RASPBERRY FLAVOR

carboxymethylcellulose sodium, citric acid, edetate disodium, FD&C blue no. 1, flavors, glycerin, microcrystalline cellulose, polysorbate 80, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucrose, xanthan gum

Drug Facts

Active ingredients

Purposes

(in each 5 mL teaspoon)

Ibuprofen 100 mg

Pain reliever/Fever reducer

Pseudoephedrine HCl 15 mg

Nasal decongestant

Uses

temporarily relieves these cold, sinus and flu symptoms:

nasal and sinus congestion headache

stuffy nose sore throat

minor aches and pains fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction which may include:

hives facial swelling asthma (wheezing) shock

Stomach bleeding warning: Taking more than recommended may cause stomach bleeding.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult a doctor right away. Do not use more than 2 days or give to children under 3 years of age unless directed by a doctor.

Do not use

if the child has ever had an allergic reaction to any pain reliever, fever reducer or nasal decongestant in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

not been drinking fluids

lost a lot of fluid due to continued vomiting or diarrhea

problems or serious side effects from taking any pain reliever, fever reducer or nasal decongestant

stomach problems that last or come back, such as heartburn, upset stomach or pain diabetes

high blood pressure, heart or kidney disease or is taking a diuretic

thyroid disease ulcers bleeding problems

Ask a doctor or pharmacist before use if the child is

under a doctor's care for any serious condition taking any other drug taking a prescription

product for anticoagulation (blood-thinning) taking any other product that contains ibuprofen

or any other pain reliever/fever reducer taking any other product that contains pseudoephedrine or any other nasal decongestant

When using this product

do not use more than directed give with food or milk if stomach upset occurs

Stop use and ask a doctor if

an allergic reaction occurs. Seek medical help right away. the child does not get any relief within first day (24 hours) of treatment fever or pain or nasal congestion gets worse or lasts for more than 3 days stomach pain or upset gets worse or lasts symptoms continue or get worse redness or swelling is present in the painful area the child gets nervous, dizzy, sleepless or sleepy any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

do not give more than directed

shake well before using

find right dose on chart below. If possible, use weight to dose; otherwise use age.

if needed, repeat dose every **6 hours**

do not use more than **4 times a day**

replace original bottle cap to maintain child resistance

measure only with dosing cup provided. Dosing cup to be used with Children's Advil Cold Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Dosing Chart

Weight (lb)	Age (yr)	Dose (teaspoons)
under 24	under 2 yr	ask a doctor
24 - 47	2 - 5 yr	1 teaspoon
48 - 95	6 - 11 yr	2 teaspoons

Other information

store at room temperature 20-25°C (68-77°F)

alcohol free

read all warnings and directions before use. Keep carton.

Inactive ingredients

carboxymethylcellulose sodium, citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, microcrystalline cellulose, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum